

Unmanned Aerial Technologies for In Situ Validation of Remote Sensing Data and Retrievals at Active Volcanoes

David Pieri, Jet Propulsion Laboratory/Caltech


Jorge Andres Diaz, CICANUM, University of Costa Rica, San Jose

*2017 HyspIRI Science and Applications Workshop
Pasadena, CA—October 14-19*

© 2017. All rights reserved.
Government sponsorship acknowledged



Jet Propulsion Laboratory California Institute of Technology




Credit Roll:

Hawaii

Andres Diaz (University of Costa Rica)
 Geoff Bland (NASA GSFC/WFF)
 Matt Fladeland (NASA ARC)
 Daniela Diaz-Chaverri (University of Costa Rica)
 Lance Christensen (JPL)
 Brendan Smith (JPL, UC Merced)
 Vince Realmuto (JPL)
 Florian Schwandner (JPL/UCLA)
 Justin Linick (JPL)
 Randy Berthold (NASA ARC ret.)
 Kent Schiffer (NASA ARC)
 Steve Patterson (NASA ARC)
 Bruce Storms (NASA ARC)
 Ric Kolyer (NASA ARC)
 Bernadette Luna (NASA ARC)
 Ernesto Corrales (UCR)
 Ted Miles (NASA GSFC/WFF ret.)
 Christophe Kern (USGS)
 Tamar Elias (USGS)
 Christina Neal (USGS)
 Rhonda Loh (NPS)
 Greg Funderburk (NPS)
 Lani and Bill Petrie (Kapapala Ranch)

Italy, Costa Rica, and Beyond

Xin Xi (GSFC, formerly NASA ARC)
 Matt Johnson (NASA ARC)
 Fabrizia Buongiorno (INGV)
 Fawzi Doumaz (INGV)
 Jack Elston (BST)
 Darby Makel (MEI)
 Kevin Baines (JPL, UWisc)
 Gary Hunter (NASA GRC)
 Ira Leifer (Bubbleology Research International)



Outline of Talk

- History of NASA Volcano UAV-based efforts
 - Science objectives
 - Case studies
 - Costa Rica
 - Italy
 - Hawaii
 - Results
 - Future Activities

Small UAV Volcanology Activities supported by
NASA SMD Earth Surface and Interior Focus Area, NASA SBIR Program, INGV, UCR
David Pieri (JPL) and Andres Diaz (University of Costa Rica)
and Government, University, and Industry Team



Scope of Activities:

- Genesis: NASA (ARC-GSFC/WFF), JPL, UCR, INGV collaboration; ISRSE Congress, Stresa, Italy, 2009

Past Deployments:

- UCR-JPL Systematic deployments of aerostats/UAVs: Turrialba Volcano 2011-2016
- NASA (Ames/GSFC-Wallops)-UCR-JPL Dragon Eye employment: Turrialba Volcano, 2013
- NASA (Ames/GSFC-Wallops)-JPL-UCR/NTCR deployment of Sierra A to Costa Rica planned for 2014; deferred with loss of a/c into Arctic Ocean
- INGV-UCR-JPL deployment to La Solfatara, Italy 2015
- INGV-UCR-JPL deployment to La Solfatara/Vulcano Island 2016

Most Recent:

- NASA (Ames/GSFC-Wallops)-JPL-UCR-USGS-NPS (UofH) deployment to Kilauea Volcano 2017

Future:

- NASA (Ames/GSFC-Wallops)-JPL-UCR-Bubbleology Inc. of SIERRA B and Dragon Eye to Salton Sea Volcanic Field 2017-2018.
- JPL-Black Swift Technology-NTCR/UCR/OVSICORI deployments in Costa Rica 2017-2018
- JPL-Makel Engineering Inc-deployment of VTOL UAV to lava sites (TBD) for NASA HOTTech Program
- JPL-NASA GRC-Black Swift Technology-USGS-NPS UAV deployment to Hawaii—TBD

Objectives

- In situ validation of remote sensing derived (e.g., ASTER, ER2-based MASTER/HyTES and AVIRIS-ng) gas and aerosol retrievals using free-flying UAV-based, aerostat-based, and ground instrumentation.
- Characterize the near surface extent, distribution, constituents, and dispersion characteristics of gas and aerosol emissions, especially SO₂ hydrolysis (e.g., Kilauea Volcano, Hawaii; Volcan Turrialba, CR).
- Improve accuracy of local SO₂ and CO₂ flux estimates using in situ airborne data (also H₂S as appropriate)
- Improve approaches to statistical representation of UAV data.
- Improve knowledge of local volcanic phenomena to mitigate hazards to local residents.
- Validate and improve transport models (applications to mitigation of airborne volcanic hazards to aviation).
- Facilitate instrument development, especially miniaturization (planetary mission testbeds).

Turrialba Volcano, Costa Rica

UAV and Aerostat deployments

High Altitude (9-13Kft ASL)

Tethered balloons



UCR VANTAR UAV


NASA DragonEye UAV

JPL/UCR SO₂ and CO₂ sensors

Dragon Eye and Aerostat Deployment to Turrialba Volcano in Costa Rica in Support of ASTER SO₂ Data Product Validation

Compact Sulfur Dioxide sensor package for Dragon Eye small UAVs and Aerostats





Measuring the spatial/temporal rate of SO₂ Hydrolysis In Situ:


- An example of in situ data from a single UAV flight—40 deployments have been made over a 2 yr. period (
- Drop-off in SO₂ vs distance from vent is visible (**dotted line**)
- Drop-off in SO₂ vs relative humidity is visible (**dotted line**)
- Range of relative humidity with distance is mostly constant
- Dominant chemical pathway in clouds/fog:



[n.b., H₂SO₄ is possibly visible in HyTES data]

- Important problem for ecology and volcanology: what are the hydrolysis rate and production constants?

SO₂ concentration data acquired along UAV flight path over Turrialba Volcano, Costa Rica




[SO₂] vs. distance from vent vs relative humidity

Constraining the sulfur dioxide degassing flux from Turrialba volcano, Costa Rica using unmanned aerial system measurements


Xin Xi, Matthew S. Johnson, Seongeun Jeong, Matthew Fladeland, David Pieri, Jorge Andres Diaz, and Geoff Bland
(*Journal of Volcanology & Geothermal Research*, 2016)

UAV data fed into reverse dispersion models generate SO₂ flux (kt/day); results consistent with observations.



Four reverse models

Measured flux range from ground stations since 2008—reasonable agreement



~20ppmv max

SO₂



La Solfatara Crater and Fumaroles, Greater Naples, Italy

UAV and Aerostat deployments

Low altitude (<500ft ASL)


INGV DJI Quadcopter—SO₂, CO₂, H₂S

UCR/JPL Mini-gas sensing package


Field Area—La Solfatarara: CO_2 , H_2S , (SO_2 , He?); “Resurgent Caldron”; now undergoing uplift and seismic activity (e.g., 31Oct14; Magnitude 2 earthquake)




Deployed a number of sensors at La Solfatara Crater, Italy on 30-31 Oct 2014



- Proof-of-concept test of new micro-miniaturization:
- Mini-gas sensor (SO_2 , CO_2 , H_2S , T, P, $\%\text{H}_2\text{O}$, GPS, and telemetry)
- Mini-mass spectrometer (250amu mass range)
- FLIR ONE iPhone-based 8-14 μm mini-imager
- Phantom Quad-copter UAV for Mini-gas lift into active fumarole.



Near Ground CO₂ Concentrations – La Solfatara, ITALY



UAV SOUNDING

Solfatara Volcano, Napoles, Italy.

Launch site: 40,82734°N 14,14149°W 101 m AMSL

Time: 1240 - 1249 UTC / October 30, 2014

Platform used: Quadcopter with Dragon_MiniGas α

Maximum altitude: 128 m AMSL

Max. CO₂ concentration: 7423 ppm @ 110.2 m AMSL

Sounding duration: 9 minutes




jorge.andres.diaz@gmail.com

gaslab.ucr@gmail.com

www.youtube.com/GasLabUCR

Near Ground H_2S Concentrations – La Solfatara, ITALY



UAV SOUNDING [PRELIMINARY DATA]

Solfatara Volcano, Naples, Italy.

Launch site: 40,82734°N 14,14149°W 101 m AMSL

Time: 1240 - 1249 UTC / October 30, 2014

Platform used: Quadcopter with Dragon_MiniGas α

Maximum altitude: 128 m AMSL

Max. H_2S concentration: 20,3 ppm @ 110.4 m AMSL

Sounding duration: 9 minutes



jorge.andres.diaz@gmail.com
gaslab.ucr@gmail.com
www.youtube.com/GasLabUCR





Vulcano Island, Tyrrhenian Sea, Italy

UAV and Aerostat deployments

Low altitude (<1000ft ASL)

UCR ItalDrone Octocopter

UCR/JPL Mini-gas package (SO_2 , CO_2 , H_2S)



Jet Propulsion Laboratory
California Institute of Technology



ItalDrone
10kg payload
With Mini-Gas sensor
Vulcano Island
Italy
September 2015
(with Andres Diaz, UCR)




Kilauea Volcano, Hawaii, USA


HyspIRI, MASTER and ASTER support

UAV and Aerostat deployments

Medium altitude (<5000ft ASL)








Boundaries of Hawaii
Volcanoes National Park


Google Earth


Data LDEO, Columbia, NSF, NOAA
© 2016 Google
Data SIO, NOAA, U.S. Navy, NGA, GEBCO
Image Landsat / Copernicus

[Tour Guide](#)

lat 20.275571° lon -156.587641° elev -11104 ft eye alt 156.00 mi









Volcanic gas and steam
plume
Halemaumau Crater,
Kilauea Volcano, Hawaii
February 12, 2017






Dragon Eye in Flight









Kilauea Panorama





Dragon Eye UAV Takeoff at Kilauea
























(From Vince Realmuto, JPL)


Brightness Temperature Difference (BTD) Time Series


- True-Color composite of visible (RGB) data. MASTER data from 2017-02-21, collected between 20:18 - 20:29 UTC.
- False-Color composite of TIR data. SO₂ plume appears in yellow.
- Brightness Temperature Difference (BTD), calculated as difference between the BT in Channel 43 (8.7 μ m) and the maximum BT over all of the MASTER TIR channels.
- BTD Values between 2017-01-19 and 2017-02-23, corresponding to the sample location marked by the white circle in Panels a – c. The decrease in (absolute) BTD for this location suggests a decrease in the concentration of SO₂ in the plume over this time interval.





(d)

STAR = Approximate location of far-field UAV deployment at Kapapala Ranch, 21 February 2017, approximate 14mi (~23km) from vent; yellow = SO₂ plume (D-stretch of MASTER ER2 Data)









Kapapala Ranch UAV test site near Pahala, HI

Panorama



Dragon Eye UAV
Takeoff at
Kapapala Ranch



Dragon Eye UAV
Landing at
Kapapala Ranch



Results

1. **NASA Dragon Eye UAVs successfully acquired airborne SO₂ and CO₂ concentration data,** and ambient atmospheric profile data between the ground and approximately 1200ft AGL for emissions from the active Kilauea summit vent during the period 12-21 Feb 2017. Ground-based SO₂ and CO₂ data were also acquired along the rim of Halemaumau Crater.
2. **Nine successful flights were accomplished.** One flight terminated with the total destruction of the aircraft and payload estimated to be within the active Kilauea lava lake. Four flights were accomplished at Kilauea Summit; five flights were accomplished over Kapapala Ranch, 23km distant from the summit, along the lower slopes of Mauna Loa.
3. **Near-field SO₂ levels ranged up to 250ppmv within the summit plume; CO₂ levels ranged up to 500ppmv within the summit plume** (approximately 100ppmv above ambient background values). Ambient non-plume UAV CO₂ slightly elevated (+10ppm) vs NOAA values (diffuse volcano emissions?).
4. **Far-field observations captured SO₂ concentrations in the range of 0.1-1.0ppmv** at altitudes up to 1500ft AGL with ambient atmospheric profile data. These values are closely consistent with Hawaii State air quality predictions for the operational area.
5. **Comparisons with ASTER and MASTER data are ongoing.** Due to operational restrictions, only one MASTER-UAV simultaneous data set was acquired in the far-field. UAV-based data were acquired during the ASTER overpass on 12 February.
6. Analysis is ongoing, especially **comparisons of UAV in situ data to the remote sensing data, as well as reverse flux modeling** (e.g., Xi et al., 2016, JVGR—Turrialba Volcano, CR)

Lessons learned from our team activities

1. Expect losses--PLAN for them! (1 of 6 DragonEyes lost to lava lake)

2. Small UAV deployments are often less expensive and more flexible than manned a/c deployments.

3. ALWAYS have telemetry with sUAVs.

4. Need all weather, day/night IFR (beyond line-of-sight) capabilities and permissions for hazard response.

5. Cutting edged government sUAV programs need to be sustainable, for advanced applications beyond the marketplace.

6. Small instrument development needs fostered/sustained.

7. VTOL /fixed-wing capabilities need development

8. Aerostats should be included in the mix.

9. Science UAV uses need advocacy!


Our JPL, NASA, USGS, US National Park Service, Univ. of Costa Rica deployment team at Kilauea,
with an Aerovironment Dragon Eye unmanned vehicle with SO₂ sensor on-board:
Volcanic Emissions Retrieval Experiment (VEREX)



Future Activities

Future UAV Deployment to the Salton Sea, CA




Salton Sea Geothermal Field
Deployment with
SIERRA-B & Dragon Eye UAVs
(Sept 2018)



Two views of the UAV-MX XPF3 (UCR) integrated within the SIERRA_B UAV airframe at NASA ARC.




Area (inside orange polygon) within which we plan to conduct flight operations with both the SIERRA and Dragon Eye UAVs.




“Gryphons” or “mudpots” (small mud volcanoes) are ubiquitous in the Salton Sea Geothermal Field, most active emitting a variety of gases: **SO₂** **CO₂** **NO_x** **CH₄** **H₂S** **OCS** **NH₃** WOW!

Mini-mass spectrometers (University of Costa Rica—GasLab)



- 100amu mass range, 3kg instrument mass;
- NASA SIERRA, UCR VANTAR (shown), or INGV RAVEN UAV implementations
- Good for lighter gases
- Prof. Andres Diaz, UCR



AIRCRAFT

SUPERSWIFT XT

- Fully autonomous operations
- Modular field swappable payload
- Electric Propulsion
- 120 minute flight time
- Up to 2.3kg payload
- NASA SBIR Project
- *Flight ceiling over 20,000ft*
- 40 mph cruise speed
- *Sealed against outside particulates*
- Laser altimeter for precision landing system and *low altitude terrain following*
- Certified under multiple NASA flight safety reviews



Future UAV, In development—USA


JPL (Pieri) Concept for Gas Chromatograph-Mass Spectrometer on UAV tether over lava lake



FY17-19 NASA HOTTech funding for ambient **Hi-Temp (500°C) electronics & sensors**
(Makel Engineering, Inc.; NASA GRC; JPL)—Hawaii? Venus?



Jet Propulsion Laboratory
California Institute of Technology



Many Thanks!

¡Muchas Gracias!

Mille Grazie!

Mahalo nui loa!

